Pain 137 (2008) 700–704

PAIN

www.elsevier.com/locate/pain

Author Index to Volume 137 (2008)

Aasvang, E.K., Brandsborg, B., Christensen, B., Jensen, T.S., Kehlet, H., Neurophysiological characterization of postherniotomy pain, **137** 173
Abernethy, A.P., Pain and sleep: Establishing bi-directional association in a population-based sample, **137** 1
Abrahamsson, C., see Lindström, E., **137** 295
Acher, F., see Goudet, C., **137** 112
Aegerter, P., see Fletcher, D., **137** 441
Affleck, G., see Keefe, F.J., **137** 164
Afra, D., see Michalsen, A., **137** 452
Alier, K.A., Chen, Y., Sollenberg, U.E., Langel, Ü., Smith, P.A., Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137** 138
Alloui, A., see Goudet, C., **137** 112
Almeida, D.M., see Edwards, R.R., **137** 202
Anderson, T., see Keefe, F.J., **137** 164
Arsenault, B., see Truchon, M., **137** 564
Asada, M., see Sakai, A., **137** 378
Ayesh, E.E., Jensen, T.S., Svensson, P., Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137** 286

Baber, K.F., see Walker, L.S., **137** 266
Baecker, M., see Michalsen, A., **137** 452
Baeyens, J.M., see Nieto, F.R., **137** 520
Baron, R., see Freynhagen, R., **137** 230
Bastuji, H., Perchet, C., Legrain, V., Montes, C., Garcia-Larrea, L., Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137** 589
Baucom, D., see Porter, L.S., **137** 306
Bendtsen, L., see Buchgreitz, L., **137** 623
Bennett, G.J., see Xiao, W.H., **137** 218
Bennett, G.J., see Xanthos, D.N., **137** 640
Bernard, R., see Van Steenwinckel, J., **137** 66
Berrang, J., see Gottschling, S., **137** 405
Blackshaw, L.A., see Lindström, E., **137** 295
Blumberg, P.M., see Zhu, H., **137** 574
Blyth, F.M., Chronic pain – Is it a public health problem? **137** 465
Borsani, E., see Martucci, C., **137** 81
Bouma, P.A.D., see Dieleman, J.P., **137** 681
Bourgoin, S., see Van Steenwinckel, J., **137** 66
Boye, M.E., see Jensen, M.P., **137** 422
Bradley, M.M., Silakowski, T., Lang, P.J., Fear of pain and defensive activation, **137** 156
Brandsborg, B., see Aasvang, E.K., **137** 173
Breivik, H., Stubhaug, A., Management of acute postoperative pain: Still a long way to go! **137** 233
Brierley, S.M., see Lindström, E., **137** 295
Brisorgueil, M.-J., see Van Steenwinckel, J., **137** 66
Brusberg, M., see Lindström, E., **137** 295
Buchgreitz, L., Lyngberg, A.C., Bendtsen, L., Jensen, R., Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137** 623
Burke, L.B., see Turk, D.C., **137** 276
Burnes, L.A., see Ikeuchi, M., **137** 662
Bussmann, J.B.J., see Schasfoort, F.C., **137** 323

Caldwell, D., see Keefe, F.J., **137** 164
Carter, E., see Webster, L., **137** 428
Casey, C.Y., Greenberg, M., Nicassio, P., Harpin, R.E., Hubbard, D., A risk factor conceptualization of pain beliefs: Response to McDonald, **137** 461
Cella, D., see Turk, D.C., **137** 276
Cendán, C.M., see Nieto, F.R., **137** 520
Cesur, Ö., see Michalsen, A., **137** 452
Chapuy, E., see Goudet, C., **137** 112
Chen, Y., see Alier, K.A., **137** 138
Chi, Y.-N., see Jiang, Y.-Q., **137** 495
Chiang, L.W., see Levin, M.E., **137** 182
Chimello, J.T., see Gaspardo, C.M., **137** 16
Cho, H., see Park, S.P., **137** 208
Christensen, B., see Aasvang, E.K., **137** 173
Chu, Y.-C., Guan, Y., Skinner, J., Raja, S.N., Johns, R.A., Tao, Y.-X., Corrigendum to "Effect of genetic knockout or pharmacologic inhibition of neuronal nitric oxide synthase on complete Freund's adjuvant-induced persistent pain" [Pain 119 (2005) 113–123], **137** 463
Clark, D.J., see Sabsovich, I., **137** 507
Cleeland, C.S., see Turk, D.C., **137** 276
Clinch, J., see Jordan, A., **137** 478
Coderre, T.J., see Xanthos, D.N., **137** 640
Colleoni, M., see Martucci, C., **137** 81
Connell, H., see Jordan, A., **137** 478
Connelly, M., see Keefe, F.J., **137** 164
Conrath, M., see Van Steenwinckel, J., **137** 66
Costa, B., see Martucci, C., **137** 81
Coste, J., Voisin, D.L., Miraucourt, L.S., Dallel, R., Luccarini, P., Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137** 340
Côté, D., see Truchon, M., **137** 564
Cowan, P., see Turk, D.C., **137** 276
Croft, P., see Jordan, K.P., **137** 652
Croft, P., see Jordan, K.P., **137** 469
Crombez, G., Eccleston, C., Van Hamme, G., De Vlieger, P., Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137** 556
Crombez, G., Eccleston, C., De Vlieger, P., Van Damme, S., De Clercq, A., Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137** 631
Cuadrado, A., see Rosa, A.O., **137** 332

doi:10.1016/S0304-3959(08)00342-4

Pain 137 (2008) 700–704

PAIN

www.elsevier.com/locate/pain

Author Index to Volume 137 (2008)

Aasvang, E.K., Brandsborg, B., Christensen, B., Jensen, T.S., Kehlet, H., Neurophysiological characterization of postherniotomy pain, **137** 173
Abernethy, A.P., Pain and sleep: Establishing bi-directional association in a population-based sample, **137** 1
Abrahamsson, C., see Lindström, E., **137** 295
Acher, F., see Goudet, C., **137** 112
Aegerter, P., see Fletcher, D., **137** 441
Affleck, G., see Keefe, F.J., **137** 164
Afra, D., see Michalsen, A., **137** 452
Alier, K.A., Chen, Y., Sollenberg, U.E., Langel, Ü., Smith, P.A., Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137** 138
Alloui, A., see Goudet, C., **137** 112
Almeida, D.M., see Edwards, R.R., **137** 202
Anderson, T., see Keefe, F.J., **137** 164
Arsenault, B., see Truchon, M., **137** 564
Asada, M., see Sakai, A., **137** 378
Ayesh, E.E., Jensen, T.S., Svensson, P., Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137** 286

Baber, K.F., see Walker, L.S., **137** 266
Baecker, M., see Michalsen, A., **137** 452
Baeyens, J.M., see Nieto, F.R., **137** 520
Baron, R., see Freynhagen, R., **137** 230
Bastuji, H., Perchet, C., Legrain, V., Montes, C., Garcia-Larrea, L., Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137** 589
Baucom, D., see Porter, L.S., **137** 306
Bendtsen, L., see Buchgreitz, L., **137** 623
Bennett, G.J., see Xiao, W.H., **137** 218
Bennett, G.J., see Xanthos, D.N., **137** 640
Bernard, R., see Van Steenwinckel, J., **137** 66
Berrang, J., see Gottschling, S., **137** 405
Blackshaw, L.A., see Lindström, E., **137** 295
Blumberg, P.M., see Zhu, H., **137** 574
Blyth, F.M., Chronic pain – Is it a public health problem? **137** 465
Borsani, E., see Martucci, C., **137** 81
Bouma, P.A.D., see Dieleman, J.P., **137** 681
Bourgoin, S., see Van Steenwinckel, J., **137** 66
Boye, M.E., see Jensen, M.P., **137** 422
Bradley, M.M., Silakowski, T., Lang, P.J., Fear of pain and defensive activation, **137** 156
Brandsborg, B., see Aasvang, E.K., **137** 173
Breivik, H., Stubhaug, A., Management of acute postoperative pain: Still a long way to go! **137** 233
Brierley, S.M., see Lindström, E., **137** 295
Brisorgueil, M.-J., see Van Steenwinckel, J., **137** 66
Brusberg, M., see Lindström, E., **137** 295
Buchgreitz, L., Lyngberg, A.C., Bendtsen, L., Jensen, R., Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137** 623
Burke, L.B., see Turk, D.C., **137** 276
Burnes, L.A., see Ikeuchi, M., **137** 662
Bussmann, J.B.J., see Schasfoort, F.C., **137** 323

Caldwell, D., see Keefe, F.J., **137** 164
Carter, E., see Webster, L., **137** 428
Casey, C.Y., Greenberg, M., Nicassio, P., Harpin, R.E., Hubbard, D., A risk factor conceptualization of pain beliefs: Response to McDonald, **137** 461
Cella, D., see Turk, D.C., **137** 276
Cendán, C.M., see Nieto, F.R., **137** 520
Cesur, Ö., see Michalsen, A., **137** 452
Chapuy, E., see Goudet, C., **137** 112
Chen, Y., see Alier, K.A., **137** 138
Chi, Y.-N., see Jiang, Y.-Q., **137** 495
Chiang, L.W., see Levin, M.E., **137** 182
Chimello, J.T., see Gaspardo, C.M., **137** 16
Cho, H., see Park, S.P., **137** 208
Christensen, B., see Aasvang, E.K., **137** 173
Chu, Y.-C., Guan, Y., Skinner, J., Raja, S.N., Johns, R.A., Tao, Y.-X., Corrigendum to "Effect of genetic knockout or pharmacologic inhibition of neuronal nitric oxide synthase on complete Freund's adjuvant-induced persistent pain" [Pain 119 (2005) 113–123], **137** 463
Clark, D.J., see Sabsovich, I., **137** 507
Cleeland, C.S., see Turk, D.C., **137** 276
Clinch, J., see Jordan, A., **137** 478
Coderre, T.J., see Xanthos, D.N., **137** 640
Colleoni, M., see Martucci, C., **137** 81
Connell, H., see Jordan, A., **137** 478
Connelly, M., see Keefe, F.J., **137** 164
Conrath, M., see Van Steenwinckel, J., **137** 66
Costa, B., see Martucci, C., **137** 81
Coste, J., Voisin, D.L., Miraucourt, L.S., Dallel, R., Luccarini, P., Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137** 340
Côté, D., see Truchon, M., **137** 564
Cowan, P., see Turk, D.C., **137** 276
Croft, P., see Jordan, K.P., **137** 652
Croft, P., see Jordan, K.P., **137** 469
Crombez, G., Eccleston, C., Van Hamme, G., De Vlieger, P., Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137** 556
Crombez, G., Eccleston, C., De Vlieger, P., Van Damme, S., De Clercq, A., Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137** 631
Cuadrado, A., see Rosa, A.O., **137** 332

doi:10.1016/S0304-3959(08)00342-4

Dahlgren, M.N., see Intondi, A.B., **137** 352
Dallel, R., see Coste, J., **137** 340
D'Eon, J.L., see Harris, C.A., **137** 609
De Clercq, A., see Crombez, G., **137** 631
De Vlieger, P., see Crombez, G., **137** 556
De Vlieger, P., see Crombez, G., **137** 631
Del Pozo, E., see Nieto, F.P., **137** 520
Derbyshire, S.W.G., see Kirwilliam, S.S., **137** 60
Dieleman, J.P., Kerklaan, J., Huygen, F.J.P.M., Bouma, P.A.D., Sturkenboom, M.C.J.M., ncidence rates and treatment of neuropathic pain conditions in the general population, **137** 681
Dionne, C., see Truchon, M., **137** 564
Distler, L., see Gottschling, S., **137** 405
Dobos, G.J., see Michalsen, A., **137** 452
Dworkin, R.H., see Turk, D.C., **137** 276
Dworkin, R.H., see O'Connor, A.B., **137** 96

Eccleston, C., see Crombez, G., **137** 631
Eccleston, C., see Crombez, G., **137** 556
Eccleston, C., see Jordan, A., **137** 478
Edwards, L., Inui, K., Ring, C., Wang, X., Kakigi, R., Pain-related evoked potentials are modulated across the cardiac cycle, **137** 488
Edwards, R.R., Almeida, D.M., Klick, B., Haythornthwaite, J.A., Smith, M.T., Duration of sleep contributes to next-day pain report in the general population, **137** 202
Egea, J., see Rosa, A.O., **137** 332
Eilers, M.A., see Intondi, A.B., **137** 352
El-Metwally, A., see Ståhl, M., **137** 316
Entrena, J.M., see Nieto, F.R., **137** 520
Ernst, E., Born to suck – The return of the leech? **137** 235
Eschalier, A., see Goudet, C., **137** 112

Farrar, J.T., see Turk, D.C., **137** 276
Fermanian, C., see Fletcher, D., **137** 441
Fillion, L., see Truchon, M., **137** 564
Fink, M., see Michalsen, A., **137** 452
Fischer, J., see Van Steenwinckel, J., **137** 66
Fletcher, D., Fermanian, C., Mardaye, A., Aegerter, P., Pain and regional anesthesia committee of the French Anesthesia and Intensive Care Society (SFAR), A patient-based national survey on postoperative pain management in France reveals significant achievements and persistent challenges, **137** 441
Flor, H., see Hermann, C., **137** 689
Formanoy, M.A.G., see Schasfoort, F.C., **137** 323
France, C.R., see Trost, Z., **137** 26
Franchi, S., see Martucci, C., **137** 81
Freynhagen, R., Rolke, R., Rutjes, A.-K., Baron, R., Tölle, T.R., Schu, S., Treede, R.-D., Pseudoradicular and radicular low-back pain – A disease continuum rather than different entities? Rebuttal: Reply to the letter "Cheese and Chalk? Missing the real anatomy" by Breck McKay, **137** 229

Gandevia, S.C., see Gustin, S.M., **137** 237
Garber, J., see Walker, L.S., **137** 266
Garcia-Larrea, L., see Bastuji, H., **137** 589
Garst, J., see Porter, L.S., **137** 306
Gaspardo, C.M., Miyase, C.I., Chimello, J.T., Martinez, F.E., Martins Linhares, M.B., Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137** 16
Geis, C., see Sabsovich, I., **137** 507
Gingrich, J.A., see Van Steenwinckel, J., **137** 66
Gortner, L., see Gottschling, S., **137** 405
Gottschling, S., Meyer, S., Gribova, I., Distler, L., Berrang, J., Gortner, L., Graf, N., Shamdeen, M.G., Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137** 405
Goudet, C., Chapuy, E., Alloui, A., Acher, F., Pin, J.-P., Eschalier, A., Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137** 112
Graf, N., see Gottschling, S., **137** 405
Greenberg, M., see Casey, C.Y., **137** 461
Greenspan, J.D., see Quiton, R.L., **137** 245
Gribova, I., see Gottschling, S., **137** 405
Guan, Y., see Chu, Y.-C., **137** 463
Gündel, H., Valet, M., Sorg, C., Huber, D., Zimmer, C., Sprenger, T., Tölle, T.R., Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137** 413
Guo, T.-Z., see Sabsovich, I., **137** 507
Gustin, S.M., Wrigley, P.J., Gandevia, S.C., Middleton, J.W., Henderson, L.A., Siddall, P.J., Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137** 237

Häkkinen, A., see Ståhl, M., **137** 316
Halder, S.L., see Nicholl, B.I., **137** 147
Hamon, M., see Van Steenwinckel, J., **137** 66
Han, J., see Zhu, H., **137** 574
Han, J.-S., see Jiang, Y.-Q., **137** 495
Han, Y., see Zhu, H., **137** 574
Handwerker, H., see Namer, B., **137** 41
Harding, G., see Turk, D.C., **137** 276
Harpin, R.E., see Casey, C.Y., **137** 461
Harris, C.A., D'Eon, J.L., Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137** 609
Havouis, R., see Rivat, C., **137** 125
Haythornthwaite, J.A., see Edwards, R.R., **137** 202
He, Q., see Zhu, H., **137** 574
Henderson, L.A., see Gustin, S.M., **137** 237
Hermann, C., Zohsel, K., Hohmeister, J., Flor, H., Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137** 689
Herrmann, D.N., see O'Connor, A.B., **137** 96
Hertz, S., see Turk, D.C., **137** 276
Hilliges, M., see Namer, B., **137** 41
Hohmeister, J., see Hermann, C., **137** 689
Hoogstraten, J., see Versloot, J., **137** 389
Hubbard, D., see Casey, C.Y., **137** 461
Huber, D., see Gündel, H., **137** 413
Hughes, P.A., see Lindström, E., **137** 295
Hussain, S., see Morley, S., **137** 670
Huygen, F.J.P.M., see Dieleman, J.P., **137** 681

Ikeuchi, M., Kolker, S.J., Burnes, L.A., Walder, R.Y., Sluka, K.A., Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137** 662
Intondi, A.B., Dahlgren, M.N., Eilers, M.A., Taylor, B.K., Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137** 352
Inui, K., see Edwards, L., **137** 488
Irving, G., see Webster, L., **137** 428

Jänig, W., What is the mechanism underlying treatment of pain by systemic application of lidocaine? **137** 5
Jansen, J.P., see Webster, L., **137** 428
Jensen, M.P., Mardekian, J., Lakshminarayanan, M., Boye, M.E., Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137** 422
Jensen, R., see Buchgreitz, L., **137** 623
Jensen, T.S., see Ayesh, E.E., **137** 286
Jensen, T.S., see Aasvang, E.K., **137** 173
Ji, R.-R., see Levin, M.E., **137** 182
Ji, Y., see Tang, B., **137** 540

Jiang, Y.-Q., Xing, G.-G., Wang, S.-L., Tu, H.-Y., Chi, Y.-N., Li, J., Liu, F.-Y., Han, J.-S., Wan, Y., Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137** 495
Jimmieson, N.L., see Johnston, V., **137** 257
Jin, J.G., see Levin, M.E., **137** 182
Johns, R.A., see Chu, Y.-C., **137** 463
Johnston, V., Jimmieson, N.L., Jull, G., Souvlis, T., Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability, **137** 257
Jordan, A., Eccleston, C., McCracken, L.M., Connell, H., Clinch, J., The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137** 478
Jordan, K.P., Thomas, E., Peat, G., Wilkie, R., Croft, P., Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137** 652
Jordan, K.P., Croft, P., Opportunities and limitations of general practice databases in pain research, **137** 469
Jull, G., see Johnston, V., **137** 257

Kakigi, R., see Edwards, L., **137** 488
Kautiainen, H., see Ståhl, M., **137** 316
Kavussanu, M., see McIntyre, D., **137** 550
Keefe, F.J., Anderson, T., Lumley, M., Caldwell, D., Stainbrook, D., Mckee, D., Waters, S.J., Connelly, M., Affleck, G., Pope, M.S., Weiss, M., Riordan, P.A., Uhlin, B.D., A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137** 164
Keefe, F.J., see Porter, L.S., **137** 306
Kehlet, H., see Aasvang, E.K., **137** 173
Kerklaan, J., see Dieleman, J.P., **137** 681
Kim, B.M., see Park, S.P., **137** 208
Kim, M., see Park, S.P., **137** 208
Kingery, W.S., see Sabsovich, I., **137** 507
Kirwilliam, S.S., Derbyshire, S.W.G., Increased bias to report heat or pain following emotional priming of pain-related fear, **137** 60
Kleoudis, C., see Webster, L., **137** 428
Klick, B., see Edwards, R.R., **137** 202
Kolker, S.J., see Ikeuchi, M., **137** 662
Koo, J.Y., see Park, S.P., **137** 208
Kumpusalo, E., see Mäntyselkä, P., **137** 34

Laboureyras, E., see Rivat, C., **137** 125
Lakshminarayanan, M., see Jensen, M.P., **137** 422
Lang, P.J., see Bradley, M.M., **137** 156
Langel, Ü., see Alier, K.A., **137** 138
Larsson, H., see Lindström, E., **137** 295
Lasko, B., see Webster, L., **137** 428
Laulin, J.-P., see Rivat, C., **137** 125
Lavery, D.J., see Levin, M.E., **137** 182
LeComte, M.D., see Petersen, K.L., **137** 395
Lee, C.H., see Park, S.P., **137** 208
Legrain, V., see Bastuji, H., **137** 589
Levin, M.E., Jin, J.G., Ji, R.-R., Tong, J., Pomonis, J.D., Lavery, D.J., Miller, S.W., Chiang, L.W., Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137** 182
Lewin, G.R., Stretching it for pain, **137** 3
Li, J., see Jiang, Y.-Q., **137** 495
Li, X., see Sabsovich, I., **137** 507
Li, Y., see Zhu, H., **137** 574
Lindström, E., Brusberg, M., Hughes, P.A., Martin, C.M., Brierley, S.M., Phillis, B.D., Martinsson, R., Abrahamsson, C., Larsson, H., Martinez, V., Blackshaw, L.A., Involvement of metabotropic glutamate 5 receptor in visceral pain, **137** 295
Liu, F.-Y., see Jiang, Y.-Q., **137** 495
Loeser, J.D., Treede, R.-D., The Kyoto protocol of IASP Basic Pain Terminology, **137** 473
López, M.G., see Rosa, A.O., **137** 332
Lorrio, S., see Rosa, A.O., **137** 332
Luccarini, P., see Coste, J., **137** 340
Lüdtke, R., see Michalsen, A., **137** 452
Lumley, M., see Keefe, F.J., **137** 164
Lyngberg, A.C., see Buchgreitz, L., **137** 623

MacDonald, M.R., The road to chronic pain disability: Negative biases or accurate pain perception? **137** 460
Macfarlane, G.J., see Nicholl, B.I. **137** 147
Magnaghi, V., see Martucci, C., **137** 81
Mäntyselkä, P., Miettola, J., Niskanen, L., Kumpusalo, E., Chronic pain, impaired glucose tolerance and diabetes: A community-based study, **137** 34
Mardaye, A., see Fletcher, D., **137** 441
Mardekian, J., see Jensen, M.P., **137** 422
Markman, J.D., see O'Connor, A.B., **137** 96
Martin, C.M., see Lindström, E., **137** 295
Martinez, F.E., see Gaspardo, C.M., **137** 16
Martinez, V., see Lindström, E., **137** 295
Martins Linhares, M.B., see Gaspardo, C.M., **137** 16
Martinsson, R., see Lindström, E., **137** 295
Martucci, C., Trovato, A.E., Costa, B., Borsani, E., Franchi, S., Magnaghi, V., Panerai, A.E., Rodella, L.F., Valsecchi, A.E., Sacerdote, P., Colleoni, M., The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137** 81
Max, M.B., see Turk, D.C., **137** 276
May, A., Chronic pain may change the structure of the brain, **137** 7
McBeth, J., see Nicholl, B.I., **137** 147
McBride, C.M., see Porter, L.S., **137** 306
McCracken, L.M., see Jordan, A., **137** 478
McGrath, P.J., The family is the crucible, **137** 471
McIntyre, D., Kavussanu, M., Ring, C., Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans, **137** 550
McKay, A.B., Cheese and Chalk? Missing the real anatomy, **137** 229
Mckee, D., see Keefe, F.J., **137** 164
Meadoff, T., see Petersen, K.L., **137** 395
Meyer, S., see Gottschling, S., **137** 405
Michalsen, A., Lüdtke, R., Cesur, Ö., Afra, D., Musial, F., Baecker, M., Fink, M., Dobos, G.J., Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial, **137** 452
Middleton, J.W., see Gustin, S.M., **137** 237
Miettola, J., see Mäntyselkä, P., **137** 34
Mikkelsson, M., see Ståhl, M., **137** 316
Miletic, G., Miletic, V., Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137** 532
Miletic, V., see Miletic, G., **137** 532
Miller, S.W., see Levin, M.E., **137** 182
Miraucourt, L.S., see Coste, J., **137** 340
Miyase, C.I., see Gaspardo, C.M., **137** 16
Montes, C., see Bastuji, H., **137** 589
Morley, S., see Sutherland, R., **137** 366
Morley, S., Williams, A., Hussain, S., Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme, **137** 670
Morlion, B., see Webster, L., **137** 428
Mortensen, E., see Webster, L., **137** 428

Moseley, G.L., Zalucki, N.M., Wiech, K., Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137** 600
Moulinoux, J.-P., see Rivat, C., **137** 125
Musial, F., see Michalsen, A., **137** 452
Musleh, M., see Nicholl, B.I., **137** 147

Na, H.S., see Park, S.P., **137** 208
Namer, B., Hilliges, M., Ørstavik, K., Schmidt, R., Weidner, C., Torebjörk, E., Handwerker, H., Schmelz, M., Endothelin1 activates and sensitizes human C-nociceptors, **137** 41
Nicassio, P., see Casey, C.Y., **137** 461
Nicholl, B.I., Halder, S.L., Macfarlane, G.J., Thompson, D.G., O'Brien, S., Musleh, M., McBeth, J., Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137** 147
Nieto, F.R., Entrena, J.M., Cendán, C.M., Pozo, E. Del, Vela, J.M., Baeyens, J.M., Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137** 520
Niskanen, L., see Mäntyselkä, P., **137** 34
Noble, F., see Rivat, C., **137** 125

O'Brien, S., see Nicholl, B.I., **137** 147
O'Connor, A.B., Schwid, S.R., Herrmann, D.N., Markman, J.D., Dworkin, R.H., Pain associated with multiple sclerosis: Systematic review and proposed classification, **137** 96
Oh, U., see Park, S.P., **137** 208
Ørstavik, K., see Namer, B., **137** 41

Pain, regional anesthesia committee of the Fr, , see Fletcher, D., **137** 441
Panerai, A.E., see Martucci, C., **137** 81
Park, S.P., Kim, B.M., Koo, J.Y., Cho, H., Lee, C.H., Kim, M., Na, H.S., Oh, U., A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137** 208
Peat, G., see Jordan, K.P., **137** 652
Peppin, J., see Webster, L., **137** 428
Perchet, C., see Bastuji, H., **137** 589
Pertovaara, A., Wei, H., Dual influence of the striatum on neuropathic hypersensitivity, **137** 50
Peters, J.W.B., see Schasfoort, F.C., **137** 323
Peters, M.M., see Petersen, K.L., **137** 395
Petersen, K.L., Meadoff, T., Press, S., Peters, M.M., LeComte, M.D., Rowbotham, M.C., Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137** 395
Phillis, B.D., see Lindström, E., **137** 295
Pierce, A., see Webster, L., **137** 428
Pin, J.-P., see Goudet, C., **137** 112
Pomonis, J.D., see Levin, M.E., **137** 182
Pope, M.S., see Keefe, F.J., **137** 164
Porter, L.S., Keefe, F.J., Garst, J., McBride, C.M., Baucom, D., Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress, **137** 306
Press, S., see Petersen, K.L., **137** 395

Quiton, R.L., Greenspan, J.D., Across- and within-session variability of ratings of painful contact heat stimuli, **137** 245

Raja, S.N., see Chu, Y.-C., **137** 463
Rappaport, B.A., see Turk, D.C., **137** 276
Revicki, D., see Turk, D.C., **137** 276
Richebé, P., see Rivat, C., **137** 125
Ring, C., see McIntyre, D., **137** 550
Ring, C., see Edwards, L., **137** 488
Riordan, P.A., see Keefe, F.J., **137** 164
Rivat, C., Richebé, P., Laboureyras, E., Laulin, J.-P., Havouis, R., Noble, F., Moulinoux, J.-P., Simonnet, G., Polyamine deficient diet to relieve pain hypersensitivity, **137** 125
Rodella, L.F., see Martucci, C., **137** 81
Rojo, A.I., see Rosa, A.O., **137** 332
Rolke, R., see Freynhagen, R., **137** 229
Rosa, A.O., Egea, J., Lorrio, S., Rojo, A.I., Cuadrado, A., López, M.G., Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137** 332
Rowbotham, M.C., see Petersen, K.L., **137** 395
Rutjes, A.-K., see Freynhagen, R., **137** 229

Sabsovich, I., Guo, T.-Z., Wei, T., Zhao, R., Li, X., Clark, D.J., Geis, C., Sommer, C., Kingery, W.S., TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137** 507
Sacerdote, P., see Martucci, C., **137** 81
Sakai, A., Asada, M., Seno, N., Suzuki, H., Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137** 378
Salminen, J.J., see Ståhl, M., **137** 316
Schasfoort, F.C., Formanoy, M.A.G., Bussmann, J.B.J., Peters, J.W.B., Tibboel, D., Stam, H.J., Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137** 323
Schmelz, M., see Namer, B., **137** 41
Schmidt, R., see Namer, B., **137** 41
Schu, S., see Freynhagen, R., **137** 229
Schwid, S.R., see O'Connor, A.B., **137** 96
Seno, N., see Sakai, A., **137** 378
Shamdeen, M.G., see Gottschling, S., **137** 405
Siddall, P.J., see Gustin, S.M., **137** 237
Silakowski, T., see Bradley, M.M., **137** 156
Simonnet, G., see Rivat, C., **137** 125
Skinner, J., see Chu, Y.-C., **137** 463
Sluka, K.A., see Ikeuchi, M., **137** 662
Smith, C.A., see Walker, L.S., **137** 266
Smith, M.T., see Edwards, R.R., **137** 202
Smith, P.A., see Alier, K.A., **137** 138
Snidow, J., see Webster, L., **137** 428
Sollenberg, U.E., see Alier, K.A., **137** 138
Sommer, C., see Sabsovich, I., **137** 507
Sorg, C., see Gündel, H., **137** 413
Souvlis, T., see Johnston, V., **137** 257
Sprenger, T., see Gündel, H., **137** 413
Ståhl, M., Kautiainen, H., El-Metwally, A., Häkkinen, A., Ylinen, J., Salminen, J.J., Mikkelsson, M., Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137** 316
Stainbrook, D., see Keefe, F.J., **137** 164
Stam, H.J., see Schasfoort, F.C., **137** 323
Stubhaug, A., see Breivik, H., **137** 233
Sturkenboom, M.C.J.M., see Dieleman, J.P., **137** 681
Sutherland, R., Morley, S., Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137** 366
Suzuki, H., see Sakai, A., **137** 378
Svensson, P., see Ayesh, E.E., **137** 286

Tang, B., Ji, Y., Traub, R.J., Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137** 540
Tao, Y.-X., see Chu, Y.-C., **137** 463
Taylor, B.K., see Intondi, A.B., **137** 352
Thomas, E., see Jordan, K.P., **137** 652
Thomas, J.S., see Trost, Z., **137** 26

Thompson, D.G., see Nicholl, B.I., **137** 147
Tibboel, D., see Schasfoort, F.C., **137** 323
Tölle, T.R., see Freynhagen, R., **137** 229
Tölle, T.R., see Gündel, H., **137** 413
Tong, J., see Levin, M.E., **137** 182
Torebjörk, E., see Namer, B., **137** 41
Traub, R.J., see Tang, B., **137** 540
Treede, R.-D., see Loeser, J.D., **137** 473
Treede, R.-D., see Freynhagen, R., **137** 229
Trost, Z., France, C.R., Thomas, J.S., Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137** 26
Trovato, A.E., see Martucci, C., **137** 81
Truchon, M., Côté, D., Fillion, L., Arsenault, B., Dionne, C., Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137** 564
Tu, H.-Y., see Jiang, Y.-Q., **137** 495
Turk, D.C., see Turner, J.A., **137** 467
Turk, D.C., Dworkin, R.H., Revicki, D., Harding, G., Burke, L.B., Cella, D., Cleeland, C.S., Cowan, P., Farrar, J.T., Hertz, S., Max, M.B., Rappaport, B.A., Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain, **137** 276
Turner, J.A., Turk, D.C., The significance of clinical significance, **137** 467

Uhlin, B.D., see Keefe, F.J., **137** 164

Valet, M., see Gündel, H., **137** 413
Valsecchi, A.E., see Martucci, C., **137** 81
Van Damme, S., see Crombez, G., **137** 631
Van Hamme, G., see Crombez, G., **137** 556
Van Steenwinckel, J., Brisorgueil, M.-J., Fischer, J., Vergé, D., Gingrich, J.A., Bourgoin, S., Hamon, M., Bernard, R., Conrath, M., Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137** 66
Veerkamp, J.S.J., see Versloot, J., **137** 389
Vela, J.M., see Nieto, F.R., **137** 520
Vergé, D., see Van Steenwinckel, J., **137** 66
Versloot, J., Veerkamp, J.S.J., Hoogstraten, J., Children's self-reported pain at the dentist, **137** 389
Voisin, D.L., see Coste, J., **137** 340

Walder, R.Y., see Ikeuchi, M., **137** 662
Walker, L.S., Baber, K.F., Garber, J., Smith, C.A., A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137** 266
Wan, Y., see Jiang, Y.-Q., **137** 495
Wang, S.-L., see Jiang, Y.-Q., **137** 495
Wang, X., see Edwards, L., **137** 488
Wang, Y., see Zhu, H., **137** 574
Waters, S.J., see Keefe, F.J., **137** 164
Webster, L., Jansen, J.P., Peppin, J., Lasko, B., Irving, G., Morlion, B., Snidow, J., Pierce, A., Mortensen, E., Kleoudis, C., Carter, E., Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137** 428
Wei, H., see Pertovaara, A., **137** 50
Wei, T., see Sabsovich, I., **137** 507
Weidner, C., see Namer, B., **137** 41
Weiss, M., see Keefe, F.J., **137** 164
Wiech, K., see Moseley, G. L., **137** 600
Wilkie, R., see Jordan, K.P., **137** 652
Williams, A., see Morley, S., **137** 670
Wrigley, P.J., see Gustin, S.M., **137** 237

Xanthos, D.N., Bennett, G.J., Coderre, T.J., Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137** 640
Xiao, W.H., Bennett, G.J., C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137** 218
Xing, G.-G., see Jiang, Y.-Q., **137** 495

Yang, Y., see Zhu, H., **137** 574
Yin, D., see Zhu, H., **137** 574
Ylinen, J., see Ståhl, M., **137** 316

Zalucki, N.M., see Moseley, G. L., **137** 600
Zhang, H., see Zhu, H., **137** 574
Zhang, Y., see Zhu, H., **137** 574
Zhao, R., see Sabsovich, I., **137** 507
Zhao, Z., see Zhu, H., **137** 574
Zhu, H., Yang, Y., Zhang, H., Han, Y., Li, Y., Zhang, Y., Yin, D., He, Q., Zhao, Z., Blumberg, P.M., Han, J., Wang, Y., Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137** 574
Zimmer, C., see Gündel, H., **137** 413
Zohsel, K., see Hermann, C., **137** 689

Pain 137 (2008) 705–714

PAIN

www.elsevier.com/locate/pain

Subject Index to Volume 137 (2008)

Accelerometry
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137**, 323
Acceptance
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
Acid
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
Acupuncture
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Adolescent chronic pain
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Adolescents
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Adrenergic receptors
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Allodynia
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137**, 218
Allodynia
Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137**, 395
Allodynia
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352
Allodynia
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Alvimopan
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Amygdala
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Analgesic
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Antagonist
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Anti-retroviral treatment
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
Anxiety
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
[Arg^8] vasopressin
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Arterial baroreceptors
Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans, **137**, 550
Arterial baroreceptors
Pain-related evoked potentials are modulated across the cardiac cycle, **137**, 488
Arthralgia
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
Arthritis
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Assessment
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Assessment
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Assessment
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Attachment
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266

doi:10.1016/S0304-3959(08)00343-6

Attention
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Attention
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
Attention
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Axon reflex flare
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
Back pain
Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137**, 564
BDI-II
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Beck Depression Inventory-second edition
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Bias
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137**, 422
Brainstem
Chronic pain may change the structure of the brain, **137**, 7
Cardiac cycle
Pain-related evoked potentials are modulated across the cardiac cycle, **137**, 488
Cardiopulmonary baroreceptors
Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans, **137**, 550
Caregivers
Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress, **137**, 306
Carpometacarpal joint
Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial, **137**, 452
Catastrophizing
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266
Catastrophizing
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Catastrophising
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
Central sensitization
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137**, 623
C-fiber
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
c-fos
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352
Children
Children's self-reported pain at the dentist, **137**, 389
Children
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Children
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Children
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Chronic abdominal pain
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266
Chronic constriction injury
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
Chronic constriction injury (CCI)
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137**, 495
Chronic constrictive injury
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Chronic disability
Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137**, 564
Chronic pain
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Chronic pain
Chronic pain may change the structure of the brain, **137**, 7
Chronic pain
Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137**, 26
Chronic pain
Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain, **137**, 276
Chronic pain
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
Chronic pain
Neurophysiological characterization of postherniotomy pain, **137**, 173
Chronic pain
Pain associated with multiple sclerosis: Systematic review and proposed classification, **137**, 96
Chronic pain
Polyamine deficient diet to relieve pain hypersensitivity, **137**, 125
Chronic pain
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Chronic pain
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
Chronic post-ischemia pain
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Chronification
Chronic pain may change the structure of the brain, **137**, 7
Cingulate cortex
Chronic pain may change the structure of the brain, **137**, 7
Clinical effectiveness
Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme, **137**, 670
Clinical trial
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164

Clinically significant change
Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme, **137**, 670
Clinician-assisted disclosure
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Cobalt protoporphyrin
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
Cognition
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137**, 589
Cognitive behavioural therapy
Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme, **137**, 670
Cohort study
Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137**, 652
Cold allodynia
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520
Colonic afferents
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Colorectal distension
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Colorectal distention
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Complement activation
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Complex regional pain syndrome
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Complex regional pain syndrome
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137**, 600
Complex regional pain syndrome
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Confirmatory factor analysis
Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137**, 564
Confirmatory factor analysis
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Constipation
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Controllability
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Coping
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266
Coping
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Cortical reorganisation
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137**, 600
CRPS-I
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Dental anxiety
Fear of pain and defensive activation, **137**, 156
Dentistry
Children's self-reported pain at the dentist, **137**, 389
Depression
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Depression
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
Descending pain modulation
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
Diabetes mellitus
Chronic pain, impaired glucose tolerance and diabetes: A community-based study, **137**, 34
Disability
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Disclosure
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Discouragement
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Distraction
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Distress
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Domains
Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain, **137**, 276
Dopamine D2 receptor
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
Dorsal horn
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Dorsal root ganglia
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Ectopic discharge
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137**, 495
Electrophysiology
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138

EMG
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137**, 323
Emotion
Fear of pain and defensive activation, **137**, 156
Emotional disclosure
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Emotional dysregulation
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
End effects
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137**, 422
Endotheline 1
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
Epidemiology
Incidence rates and treatment of neuropathic pain conditions in the general population, **137**, 681
Epidemiology
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Estradiol
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Estrogen receptor
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Experimental pain models
Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137**, 395
Exposure
Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137**, 26
Fear
Fear of pain and defensive activation, **137**, 156
Fear
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
fMRI
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Follow-up
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137**, 623
Formalin
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352
Formalin test
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
Fos immunoreactivity
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Functioning
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
GABA
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
GABAergic inhibition
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
Gastrointestinal motility
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Gender
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Gene delivery
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137**, 574
Gene expression microarray
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
General population
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137**, 623
Generalization
Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137**, 26
Glial cell line-derived neurotrophic factor
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Glutamate
Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137**, 112
Glutamate
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Gonadal hormones
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
GsMTx4
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137**, 208
Haeme oxygenase-1
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
Headache
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137**, 623
Headache
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Health-related quality of life
Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain, **137**, 276
Health-related quality of life
Pain associated with multiple sclerosis: Systematic review and proposed classification, **137**, 96
Heat
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Heat hyperalgesia
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520

Histamine
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
HO-1
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
5-HT_{1A} receptor
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
5-HT2A receptor
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
Hyperalgesia
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137**, 218
Hyperalgesia
Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137**, 395
Hyperalgesia
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Hyperalgesia
Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137**, 112
Hyperalgesia
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352
Hyperalgesia
Polyamine deficient diet to relieve pain hypersensitivity, **137**, 125
Hyperalgesia
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
Hyperalgesia
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Hyperglycaemia
Chronic pain, impaired glucose tolerance and diabetes: A community-based study, **137**, 34
Hyperpolarization-activated cyclic nucleotide-gated cation channel
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137**, 495
Hypervigilance
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
I_h
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137**, 495
IBS
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Idiopathic chronic pain
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
IL-1β
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
IL-6
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
Imagined movements
Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137**, 237
Imaging
Chronic pain may change the structure of the brain, **137**, 7
Immune system
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Immunocytochemistry
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
Impact
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Improvement
A patient-based national survey on postoperative pain management in France reveals significant achievements and persistent challenges, **137**, 441
Incidence rate
Incidence rates and treatment of neuropathic pain conditions in the general population, **137**, 681
Inflammation
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
Inflammatory heat hypersensitivity
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137**, 574
Inflammatory pain
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137**, 218
Inflammatory pain
Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137**, 112
Inguinal hernia
Neurophysiological characterization of postherniotomy pain, **137**, 173
Intrathecal injection
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137**, 574
Inventory
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Irritable bowel syndrome
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Ischemia
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Joint
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
Ketamine
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
Kinesiophobia
Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137**, 26

Knockout mice
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
Knock-out mice
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
L-NIO
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Laser
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Laser Doppler
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Laser evoked potential
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137**, 589
Learned helplessness
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Leech therapy
Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial, **137**, 452
LEP
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137**, 589
Lidocaine
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137**, 218
Local anesthesia
Children's self-reported pain at the dentist, **137**, 389
Lung cancer
Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress, **137**, 306
Measure
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Mechanical allodynia
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520
Mechanical pain
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137**, 208
Metabotropic glutamate receptor
Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137**, 112
mGluR5
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Mice
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
MicroCT scanning
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Microneurography
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
Migraine
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Motor cortex
Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137**, 237
Multi-dimensional
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Multiple sclerosis
Pain associated with multiple sclerosis: Systematic review and proposed classification, **137**, 96
Muscle
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels, **137**, 218
Musculoskeletal
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
National survey
A patient-based national survey on postoperative pain management in France reveals significant achievements and persistent challenges, **137**, 441
NCAM
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Neck pain
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Neck pain
Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability, **137**, 257
Neuralgia
Incidence rates and treatment of neuropathic pain conditions in the general population, **137**, 681
Neurogenic pain
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Neuroimaging
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Neuroinflammation
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Neuropathic
Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137**, 237
Neuropathic pain
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137**, 208
Neuropathic pain
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat, **137**, 495
Neuropathic pain
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Neuropathic pain
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
Neuropathic pain
Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain, **137**, 112

Neuropathic pain
Incidence rates and treatment of neuropathic pain conditions in the general population, **137**, 681
Neuropathic pain
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Neuropathic pain
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
Neuropathic pain
Neurophysiological characterization of postherniotomy pain, **137**, 173
Neuropathic pain
Pain associated with multiple sclerosis: Systematic review and proposed classification, **137**, 96
Neuropathic pain
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Neuropathic pain
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520
Neuropathic pain
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
Neuropathic pain
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Neuropeptide
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Neuroplasticity
Chronic pain may change the structure of the brain, **137**, 7
Neurotrophins
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
NICU
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Nitric oxide synthase
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
NMDA
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
NMDA receptor
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Nociception
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Nociception
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137**, 340
Nociception
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
Nociception
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
NR2B phosphorylation
Polyamine deficient diet to relieve pain hypersensitivity, **137**, 125
Nrf2
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice, **137**, 332
Opioid
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Opioid
Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137**, 395
Orbitofrontal cortex
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Orofacial
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137**, 340
Osteoarthritis
Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial, **137**, 452
Paclitaxel
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520
Pain
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Pain
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Pain
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Pain
Children's self-reported pain at the dentist, **137**, 389
Pain
Chronic pain, impaired glucose tolerance and diabetes: A community-based study, **137**, 34
Pain
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Pain
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137**, 340
Pain
Duration of sleep contributes to next-day pain report in the general population, **137**, 202
Pain
Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans, **137**, 550
Pain
Fear of pain and defensive activation, **137**, 156
Pain
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631

Pain
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Pain
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137**, 589
Pain
Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137**, 237
Pain
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice, **137**, 662
Pain
Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137**, 652
Pain assessment
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137**, 422
Pain behavior
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Pain behaviour
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
Pain beliefs
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266
Pain indicators
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137**, 323
Pain ratings
Pain-related evoked potentials are modulated across the cardiac cycle, **137**, 488
Pain sensitivity
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study, **137**, 623
Pain sensitization
Polyamine deficient diet to relieve pain hypersensitivity, **137**, 125
Pain-related evoked potentials
Pain-related evoked potentials are modulated across the cardiac cycle, **137**, 488
Pain-related fear
Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task, **137**, 26
Parent behavior
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Parents
The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain, **137**, 478
Patch clamp
Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin, **137**, 138
Patient-reported outcomes
Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain, **137**, 276
Peak effects
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137**, 422
Peripheral
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain, **137**, 182
Physical therapy
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137**, 600
PKA
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Polyamine deficient diet
Polyamine deficient diet to relieve pain hypersensitivity, **137**, 125
Population-based
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Population-based study
Chronic pain, impaired glucose tolerance and diabetes: A community-based study, **137**, 34
Postherniotomy
Neurophysiological characterization of postherniotomy pain, **137**, 173
Postoperative pain
A patient-based national survey on postoperative pain management in France reveals significant achievements and persistent challenges, **137**, 441
Post-operative pain
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain, **137**, 422
Potassium-chloride co-transporters
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
PPADS
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
Predictability
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Preterm
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Pre-verbal infants
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137**, 323
Problem solving
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients, **137**, 556
Problem solving
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Prognosis
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Prospective
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Protein kinase D1 (PKD1)
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137**, 574
Psychological intervention
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164

Psychometrics
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain, **137**, 609
Psychophysiology
Fear of pain and defensive activation, **137**, 156
Psychosocial
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Psychosocial factors
Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137**, 564
Purinergic antagonism
The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice, **137**, 81
Quantitative sensory measures
Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability, **137**, 257
Quantitative sensory testing
Neurophysiological characterization of postherniotomy pain, **137**, 173
Quantitative sensory tests
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
Randomized
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Randomized controlled trial
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain, **137**, 428
Randomized trial
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Randomized trial
Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial, **137**, 452
Rat
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352
Reliable change index
Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme, **137**, 670
Repeatability
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Repeated doses
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Residence characteristics
Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137**, 652
RET
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Rheumatoid arthritis
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Risk factors
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study, **137**, 316
Risk factors
Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study, **137**, 147
Risk factors
Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137**, 652
Self-efficacy
Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress, **137**, 306
Self-discrepancy
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
Self-pain enmeshment
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain, **137**, 366
Self-report
Children's self-reported pain at the dentist, **137**, 389
Sensation
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137**, 600
Sensitization
Endothelin1 activates and sensitizes human C-nociceptors, **137**, 41
Sex
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Signal analysis
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study, **137**, 323
SIN-1
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Sleep
Duration of sleep contributes to next-day pain report in the general population, **137**, 202
Sleep
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals, **137**, 589
Social responses
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Social risks
Social risks for disabling pain in older people: A prospective study of individual and area characteristics, **137**, 652
Sodium nitroprusside
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Solicitousness
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents, **137**, 689
Somatic
Increased bias to report heat or pain following emotional priming of pain-related fear, **137**, 60
Somatoform pain disorder
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Spared nerve injury
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain, **137**, 352

Spider toxin
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137**, 208
Spinal cord
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Spinal cord
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain, **137**, 378
Spinal cord injuries
Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury, **137**, 237
Startle reflex
Fear of pain and defensive activation, **137**, 156
sTNF-R1
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Stress
A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects? **137**, 164
Striatum
Dual influence of the striatum on neuropathic hypersensitivity, **137**, 50
Structural equation modeling
Low-back-pain related disability: An integration of psychological risk factors into the stress process model, **137**, 564
Substance P
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137**, 340
Sucrose
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Symptom management
Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress, **137**, 306
Synaptic plasticity
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn, **137**, 532
Tactile acuity
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain, **137**, 600
Tarantula
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain, **137**, 208
Tension type headache
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial, **137**, 405
Tetrodotoxin
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice, **137**, 520
Tibia fracture
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
TMJ pain
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
TNF
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I, **137**, 507
Tolerance
Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers, **137**, 395
Transient receptor potential V1 (TRPV1)
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity, **137**, 574
Treatment
Incidence rates and treatment of neuropathic pain conditions in the general population, **137**, 681
Trigeminal
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons, **137**, 340
Trigeminal physiology
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients, **137**, 286
Typology
A typology of pain coping strategies in pediatric patients with chronic abdominal pain, **137**, 266
Ultrastructure
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66
Upper limb nociceptive flexion reflex
Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans, **137**, 550
Vasoconstriction
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain, **137**, 640
Ventromedial prefrontal cortex
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder, **137**, 413
Very low birth weight
Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates? **137**, 16
Visceral nociception
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Visceral pain
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Visceromotor response
Involvement of metabotropic glutamate 5 receptor in visceral pain, **137**, 295
Visceromotor response
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat, **137**, 540
Visual analog scale
Across- and within-session variability of ratings of painful contact heat stimuli, **137**, 245
Voxel-based morphometry (VBM)
Chronic pain may change the structure of the brain, **137**, 7
Worry
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain, **137**, 631
Zalcitabine
Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse, **137**, 66

Pain 137 (2008) 715–718

PAIN

www.elsevier.com/locate/pain

Contents to Volume 137 (2008)

Number 1 – 30 June 2008

EDITORIALS

Pain and sleep: Establishing bi-directional association in a population-based sample
A.P. Abernethy (USA) 1

Stretching it for pain
G.R. Lewin (Germany) 3

What is the mechanism underlying treatment of pain by systemic application of lidocaine?
W. Jänig (Germany) 5

REVIEW

Chronic pain may change the structure of the brain
A. May (Germany) 7

RESEARCH PAPERS

Is pain relief equally efficacious and free of side effects with repeated doses of oral sucrose in preterm neonates?
C.M. Gaspardo, C.I. Miyase, J.T. Chimello, F.E. Martinez, M.B. Martins Linhares (Brazil) 16

Exposure to movement in chronic back pain: Evidence of successful generalization across a reaching task
Z. Trost, C.R. France, J.S. Thomas (USA) 26

Chronic pain, impaired glucose tolerance and diabetes: A community-based study
P. Mäntyselkä, J. Miettola, L. Niskanen, E. Kumpusalo (Finland) 34

Endothelin1 activates and sensitizes human C-nociceptors
B. Namer, M. Hilliges, K. Ørstavik, R. Schmidt, C. Weidner, E. Torebjörk, H. Handwerker, M. Schmelz (Germany, Sweden, Norway) 41

Dual influence of the striatum on neuropathic hypersensitivity
A. Pertovaara, H. Wei (Finland) 50

Increased bias to report heat or pain following emotional priming of pain-related fear
S.S. Kirwilliam, S.W.G. Derbyshire (UK) 60

Role of spinal serotonin 5-HT2A receptor in 2′,3′-dideoxycytidine-induced neuropathic pain in the rat and the mouse
J. Van Steenwinckel, M.-J. Brisorgueil, J. Fischer, D. Vergé, J.A. Gingrich, S. Bourgoin, M. Hamon, R. Bernard, M. Conrath (France, USA) 66

The purinergic antagonist PPADS reduces pain related behaviours and interleukin-1β, interleukin-6, iNOS and nNOS overproduction in central and peripheral nervous system after peripheral neuropathy in mice
C. Martucci, A.E. Trovato, B. Costa, E. Borsani, S. Franchi, V. Magnaghi, A.E. Panerai, L.F. Rodella, A.E. Valsecchi, P. Sacerdote, M. Colleoni (Italy) 81

Pain associated with multiple sclerosis: Systematic review and proposed classification
A.B. O'Connor, S.R. Schwid, D.N. Herrmann, J.D. Markman, R.H. Dworkin (USA) 96

Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain
C. Goudet, E. Chapuy, A. Alloui, F. Acher, J.-P. Pin, A. Eschalier (France) 112

Polyamine deficient diet to relieve pain hypersensitivity
C. Rivat, P. Richebé, E. Laboureyras, J.-P. Laulin, R. Havouis, F. Noble, J.-P. Moulinoux, G. Simonnet (France) 125

Selective stimulation of GalR1 and GalR2 in rat *substantia gelatinosa* reveals a cellular basis for the anti- and pro-nociceptive actions of galanin
K.A. Alier, Y. Chen, U.E. Sollenberg, Ü. Langel, P.A. Smith (Canada, Sweden) 138

Psychosocial risk markers for new onset irritable bowel syndrome – Results of a large prospective population-based study
B.I. Nicholl, S.L. Halder, G.J. Macfarlane, D.G. Thompson, S. O'Brien, M. Musleh, J. McBeth (UK) 147

Fear of pain and defensive activation
M.M. Bradley, T. Silakowski, P.J. Lang (USA) 156

A randomized, controlled trial of emotional disclosure in rheumatoid arthritis: Can clinician assistance enhance the effects?
F.J. Keefe, T. Anderson, M. Lumley, D. Caldwell, D. Stainbrook, D. Mckee, S.J. Waters, M. Connelly, G. Affleck, M.S. Pope, M. Weiss, P.A. Riordan, B.D. Uhlin (USA) 164

doi:10.1016/S0304-3959(08)00344-8

Neurophysiological characterization of postherniotomy pain
E.K. Aasvang, B. Brandsborg, B. Christensen, T.S. Jensen, H. Kehlet (Denmark) 173
Complement activation in the peripheral nervous system following the spinal nerve ligation model of neuropathic pain
M.E. Levin, J.G. Jin, R.-R. Ji, J. Tong, J.D. Pomonis, D.J. Lavery, S.W. Miller, L.W. Chiang (USA, Canada) 182
Duration of sleep contributes to next-day pain report in the general population
R.R. Edwards, D.M. Almeida, B. Klick, J.A. Haythornthwaite, M.T. Smith (USA) 202
A tarantula spider toxin, GsMTx4, reduces mechanical and neuropathic pain
S.P. Park, B.M. Kim, J.Y. Koo, H. Cho, C.H. Lee, M. Kim, H.S. Na, U. Oh (Republic of Korea) 208
C-fiber spontaneous discharge evoked by chronic inflammation is suppressed by a long-term infusion of lidocaine yielding nanogram per milliliter plasma levels
W.H. Xiao, G.J. Bennett (Canada) 218

LETTERS TO THE EDITOR

Cheese and Chalk? Missing the real anatomy
A.B. McKay (Australia) 229
Pseudoradicular and radicular low-back pain – A disease continuum rather than different entities? Rebuttal: Reply to the letter "Cheese and Chalk? Missing the real anatomy" by Breck McKay
R. Freynhagen, R. Rolke, A.-K. Rutjes, R. Baron, T.R. Tölle, S. Schu, R.-D. Treede (Germany) 229

Number 2 – 15 July 2008

EDITORIALS

Management of acute postoperative pain: Still a long way to go!
H. Breivik, A. Stubhaug (Norway) 233
Born to suck – The return of the leech?
E. Ernst (UK) 235

RESEARCH PAPERS

Movement imagery increases pain in people with neuropathic pain following complete thoracic spinal cord injury
S.M. Gustin, P.J. Wrigley, S.C. Gandevia, J.W. Middleton, L.A. Henderson, P.J. Siddall (Australia) 237
Across- and within-session variability of ratings of painful contact heat stimuli
R.L. Quiton, J.D. Greenspan (USA) 245
Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability
V. Johnston, N.L. Jimmieson, G. Jull, T. Souvlis (Australia) 257
A typology of pain coping strategies in pediatric patients with chronic abdominal pain
L.S. Walker, K.F. Baber, J. Garber, C.A. Smith (USA) 266
Identifying important outcome domains for chronic pain clinical trials: An IMMPACT survey of people with pain
D.C. Turk, R.H. Dworkin, D. Revicki, G. Harding, L.B. Burke, D. Cella, C.S. Cleeland, P. Cowan, J.T. Farrar, S. Hertz, M.B. Max, B.A. Rappaport (USA) 276
Effects of intra-articular ketamine on pain and somatosensory function in temporomandibular joint arthralgia patients
E.E. Ayesh, T.S. Jensen, P. Svensson (Denmark) 286
Involvement of metabotropic glutamate 5 receptor in visceral pain
E. Lindström, M. Brusberg, P.A. Hughes, C.M. Martin, S.M. Brierley, B.D. Phillis, R. Martinsson, C. Abrahamsson, H. Larsson, V. Martinez, L.A. Blackshaw (Sweden, Australia) 295
Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms and distress
L.S. Porter, F.J. Keefe, J. Garst, C.M. McBride, D. Baucom (USA) 306
Non-specific neck pain in schoolchildren: Prognosis and risk factors for occurrence and persistence. A 4-year follow-up study
M. Ståhl, H. Kautiainen, A. El-Metwally, A. Häkkinen, J. Ylinen, J.J. Salminen, M. Mikkelsson (Finland, UK) 316
Objective and continuous measurement of peripheral motor indicators of pain in hospitalized infants: A feasibility study
F.C. Schasfoort, M.A.G. Formanoy, J.B.J. Bussmann, J.W.B. Peters, D. Tibboel, H.J. Stam (The Netherlands) 323
Nrf2-mediated haeme oxygenase-1 up-regulation induced by cobalt protoporphyrin has antinociceptive effects against inflammatory pain in the formalin test in mice
A.O. Rosa, J. Egea, S. Lorrio, A.I. Rojo, A. Cuadrado, M.G. López (Spain) 332
Dorsal horn NK1-expressing neurons control windup of downstream trigeminal nociceptive neurons
J. Coste, D.L. Voisin, L.S. Miraucourt, R. Dallel, P. Luccarini (France) 340
Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain
A.B. Intondi, M.N. Dahlgren, M.A. Eilers, B.K. Taylor (USA) 352
Self-pain enmeshment: Future possible selves, sociotropy, autonomy and adjustment to chronic pain
R. Sutherland, S. Morley (UK) 366
Involvement of neural cell adhesion molecule signaling in glial cell line-derived neurotrophic factor-induced analgesia in a rat model of neuropathic pain
A. Sakai, M. Asada, N. Seno, H. Suzuki (Japan) 378
Children's self-reported pain at the dentist
J. Versloot, J.S.J. Veerkamp, J. Hoogstraten (Canada, The Netherlands) 389

Changes in morphine analgesia and side effects during daily subcutaneous administration in healthy volunteers
K.L. Petersen, T. Meadoff, S. Press, M.M. Peters, M.D. LeComte, M.C. Rowbotham (USA) 395
Laser acupuncture in children with headache: A double-blind, randomized, bicenter, placebo-controlled trial
S. Gottschling, S. Meyer, I. Gribova, L. Distler, J. Berrang, L. Gortner, N. Graf, M.G. Shamdeen (Germany) 405
Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder
H. Gündel, M. Valet, C. Sorg, D. Huber, C. Zimmer, T. Sprenger, T.R. Tölle (Germany) 413
Validity of 24-h recall ratings of pain severity: Biasing effects of "Peak" and "End" pain
M.P. Jensen, J. Mardekian, M. Lakshminarayanan, M.E. Boye (USA) 422
Alvimopan, a peripherally acting mu-opioid receptor (PAM-OR) antagonist for the treatment of opioid-induced bowel dysfunction: Results from a randomized, double-blind, placebo-controlled, dose-finding study in subjects taking opioids for chronic non-cancer pain
L. Webster, J.P. Jansen, J. Peppin, B. Lasko, G. Irving, B. Morlion, J. Snidow, A. Pierce, E. Mortensen, C. Kleoudis, E. Carter (USA, Germany, Canada, Belgium) 428
A patient-based national survey on postoperative pain management in France reveals significant achievements and persistent challenges
D. Fletcher, C. Fermanian, A. Mardaye, P. Aegerter, Pain and regional anesthesia committee of the French Anesthesia and Intensive Care Society (SFAR) (France) 441
Effectiveness of leech therapy in women with symptomatic arthrosis of the first carpometacarpal joint: A randomized controlled trial
A. Michalsen, R. Lüdtke, Ö. Cesur, D. Afra, F. Musial, M. Baecker, M. Fink, G.J. Dobos (Germany) 452

LETTERS TO THE EDITOR

The road to chronic pain disability: Negative biases or accurate pain perception?
M.R. MacDonald (Canada) 460
A risk factor conceptualization of pain beliefs: Response to McDonald
C.Y. Casey, M. Greenberg, P. Nicassio, R.E. Harpin, D. Hubbard (USA) 461

CORRIGENDUM

Corrigendum to "Effect of genetic knockout or pharmacologic inhibition of neuronal nitric oxide synthase on complete Freund's adjuvant-induced persistent pain" [Pain 119 (2005) 113–123]
Y.-C. Chu, Y. Guan, J. Skinner, S.N. Raja, R.A. Johns, Y.-X. Tao (USA, Taiwan) 463

Number 3 – 31 July 2008

EDITORIALS

Chronic pain – Is it a public health problem?
F.M. Blyth (Australia) 465
The significance of clinical significance
J.A. Turner, D.C. Turk (USA) 467
Opportunities and limitations of general practice databases in pain research
K.P. Jordan, P. Croft (UK) 469
The family is the crucible
P.J. McGrath (Canada) 471

TOPICAL REVIEW

The Kyoto protocol of IASP Basic Pain Terminology
J.D. Loeser, R.-D. Treede (USA, Germany) 473

RESEARCH PAPERS

The Bath Adolescent Pain – Parental Impact Questionnaire (BAP-PIQ): Development and preliminary psychometric evaluation of an instrument to assess the impact of parenting an adolescent with chronic pain
A. Jordan, C. Eccleston, L.M. McCracken, H. Connell, J. Clinch (UK) 478
Pain-related evoked potentials are modulated across the cardiac cycle
L. Edwards, K. Inui, C. Ring, X. Wang, R. Kakigi (UK, Japan) 488
Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat
Y.-Q. Jiang, G.-G. Xing, S.-L. Wang, H.-Y. Tu, Y.-N. Chi, J. Li, F.-Y. Liu, J.-S. Han, Y. Wan (China) 495
TNF signaling contributes to the development of nociceptive sensitization in a tibia fracture model of complex regional pain syndrome type I
I. Sabsovich, T.-Z. Guo, T. Wei, R. Zhao, X. Li, D.J. Clark, C. Geis, C. Sommer, W.S. Kingery (USA, Germany) 507
Tetrodotoxin inhibits the development and expression of neuropathic pain induced by paclitaxel in mice
F.R. Nieto, J.M. Entrena, C.M. Cendán, E. Del Pozo, J.M. Vela, J.M. Baeyens (Spain) 520
Loose ligation of the sciatic nerve is associated with TrkB receptor-dependent decreases in KCC2 protein levels in the ipsilateral spinal dorsal horn
G. Miletic, V. Miletic (USA) 532
Estrogen alters spinal NMDA receptor activity via a PKA signaling pathway in a visceral pain model in the rat
B. Tang, Y. Ji, R.J. Traub (USA) 540

Effects of arterial and cardiopulmonary baroreceptor activation on the upper limb nociceptive flexion reflex and electrocutaneous pain in humans
D. McIntyre, M. Kavussanu, C. Ring (UK) 550
Attempting to solve the problem of pain: A questionnaire study in acute and chronic pain patients
G. Crombez, C. Eccleston, G. Van Hamme, P. De Vlieger (Belgium, UK) 556
Low-back-pain related disability: An integration of psychological risk factors into the stress process model
M. Truchon, D. Côté, L. Fillion, B. Arsenault, C. Dionne (Canada) 564
Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity
H. Zhu, Y. Yang, H. Zhang, Y. Han, Y. Li, Y. Zhang, D. Yin, Q. He, Z. Zhao, P.M. Blumberg, J. Han, Y. Wang (PR China, USA) 574
Laser evoked responses to painful stimulation persist during sleep and predict subsequent arousals
H. Bastuji, C. Perchet, V. Legrain, C. Montes, L. Garcia-Larrea (France, Belgium, Spain) 589
Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain
G.L. Moseley, N.M. Zalucki, K. Wiech (UK, Australia) 600
Psychometric properties of the Beck Depression Inventory-Second Edition (BDI-II) in individuals with chronic pain
C.A. Harris, J.L. D'Eon (Canada) 609
Increased pain sensitivity is not a risk factor but a consequence of frequent headache: A population-based follow-up study
L. Buchgreitz, A.C. Lyngberg, L. Bendtsen, R. Jensen (Denmark) 623
Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain
G. Crombez, C. Eccleston, P. De Vlieger, S. Van Damme, A. De Clercq (Belgium, UK) 631
Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain
D.N. Xanthos, G.J. Bennett, T.J. Coderre (Canada) 640
Social risks for disabling pain in older people: A prospective study of individual and area characteristics
K.P. Jordan, E. Thomas, G. Peat, R. Wilkie, P. Croft (UK) 652
Role of ASIC3 in the primary and secondary hyperalgesia produced by joint inflammation in mice
M. Ikeuchi, S.J. Kolker, L.A. Burnes, R.Y. Walder, K.A. Sluka (USA) 662
Estimating the clinical effectiveness of cognitive behavioural therapy in the clinic: Evaluation of a CBT informed pain management programme
S. Morley, A. Williams, S. Hussain (UK) 670
Incidence rates and treatment of neuropathic pain conditions in the general population
J.P. Dieleman, J. Kerklaan, F.J.P.M. Huygen, P.A.D. Bouma, M.C.J.M. Sturkenboom (The Netherlands) 681
Dimensions of pain-related parent behavior: Development and psychometric evaluation of a new measure for children and their parents
C. Hermann, K. Zohsel, J. Hohmeister, H. Flor (Germany) 689

Author Index to Volume 137 700

Subject Index to Volume 137 705

Contents to Volume 137 715